

FIELD THEORETICAL COMPUTATIONS OF STATIC CONDUCTIVITY FOR A WHITE NOISE MAGNETIC SYSTEM

Cyril FURTLEHNER¹

*Institute of Physics, University of Oslo
P.O. Box 1048 Blindern
N-0316 Oslo, Norway*

Abstract

We consider an electron coupled to a random magnetic field with local correlations and eventually non-zero average value. Starting from the source-term formalism of Levine, Libby and Pruisken, we define a generating function for static transport coefficients. We use a non-hermitian representation of the Hamiltonian, and a functional representation of the generating function, where the integration variable consists of non-local supermatrices, due to the non-locality of the Aharonov-Bohm interaction. The conductivity coefficients are computed using a specific saddle-point procedure, which incorporates ladder diagrams contributions. The situation with an external uniform magnetic field is considered with the two different spin projection possibilities. The limit of vanishingly small external magnetic field is unstable with respect to spin flip, because of $E = 0$ states. In particular a quantized step is obtained when comparing the Hall conductance between the two possible spin projections.

¹cyril.furtlehner@fys.uio.no

1. Introduction

The problem of an electron moving in a random magnetic field has been in the recent years subject of a great interest in particular because of its relevance to the Hall problem, especially at half filling [1]. It may be defined in many ways, either on a Lattice with random fluxes [2, 7] or in the continuum with various choices for the statistical correlations of the magnetic field, by considering for example random distribution of point-like vortices [8, 9]. The spectrum properties of these models are now well established. Although weak-localization has been studied [11], diffusion modes have been identified, and an effective field theory for weak disorder has been constructed [12], the question of localization is however still under debate. Most recent results seem to converge to the idea that all states are localized except at $E = 0$ for lattice models [13, 19]. The question of transport coefficient has been less studied. As noticed by Lusakowski and Turski [20], there might be some additional features due to the random nature of the velocity operator. Concerning this aspect, some qualitative and quantitative perturbative results have been obtained in the situation of random magnetic impurities and strong uniform magnetic field, in particular oscillations of the Hall conductance [22, 23].

In the present article, we are interested in using some field theoretical methods, originally devised in the context of localization, to consider this question of transport coefficients. Especially we would like, as a first step, to compute the SCBA approximation in a well defined manner, using this field theoretical language. We will consider a two dimensional model where the magnetic fluctuations are distributed with uncorrelated gaussian distribution and are added to an eventual uniform magnetic field. In the first part, as a preliminary, we use the source term formalism of Levine, Libby and Pruisken to define a generating function for static transport coefficients, which can be seen as a vacuum to vacuum transition amplitude at the Fermi energy, and necessitate to consider sources in matrix form. As a byproduct it is a generating function for Ward identities. In the second part, the model under consideration is specified, and a non-unitary transformation is performed, which leaves the generating function invariant and simplifies the averaging procedure. The generating function is then expressed, using functional integration over non-local replicated supermatrices, in consequence of the non-locality of the Aharonov-Bohm interaction. This is the main technical difference with the standard random scalar potential. The saddle-point is then exactly solved and leads to semi-circle type expression for the DOS. For the issue of transport coefficients, the saddle point equation is then reconsidered in presence of the sources, which define in an unambiguous manner the saddle-point expression for these transport coefficient, and allow to eventually incorporate ladder diagrams contributions, which are known to be exclusively responsible of IR divergencies in absence of time reversal symmetry. Some perspective and natural extensions of this work, will then be drawn in the conclusive section.

2. Generating function for static transport coefficients

2.1 Smrcka-Streda formula

We start with the expressions for the static conductivity coefficients, obtained from the linear response theory. Given the general form of the Hamiltonian

$$H = \frac{1}{2m}(\mathbf{p} - e\mathbf{A})^2 + V(\mathbf{r}) \quad (1)$$

and assuming a zero temperature Fermi-Dirac equilibrium distribution, longitudinal and transverse transport coefficients can be expressed as Fermi-energy quantities, in the following way

($\hbar = 1$) [24]

$$\sigma_{xx} = \frac{\pi e^2}{V} \text{Tr}[v_x \delta(E_F - H) v_x \delta(E_F - H)] \quad (2)$$

and

$$\sigma_{xy} = \sigma_{xy}^I + \sigma_{xy}^{II} \quad (3)$$

$$\sigma_{xy}^I = \frac{ie^2}{2V} \text{Tr}[v_x G^+ v_y \delta(E_F - H) - v_x \delta(E_F - H) G^- v_y] \quad (4)$$

$$\sigma_{xy}^{II} = \frac{e^2}{2V} \text{Tr}[(xv_y - yv_x) \delta(E_F - H)] \quad (5)$$

$G^\pm = (E_F - H \pm i\eta)^{-1}$ are the advanced (resp. retarded) Green function. $v_i = p_i - eA_i$ are the components of the velocity operator. V is the volume, and these expressions represent an average over the sample of local current-current correlation function, concerning σ_{xx} and σ_{xy}^I , although σ_{xy}^{II} is the magnetization at the Fermi energy per unit volume of the sample. It is already to be noticed that this last formula is not well defined unless edges are taken into account [26, 27]. A way to avoid a cumbersome regularization is to use the Streda formula [25]

$$\sigma_{xy}^{II} = e \frac{\partial N(E)}{\partial B} \Big|_{E=E_F} \quad (6)$$

where $N(E)$ is the integrated density of states. Since the model we have in mind is two dimensional, it will be convenient to use complexe notations ($z = x + iy$, $A_z = A_x + iA_y$, $\partial_z = \frac{1}{2}(\partial_x - i\partial_y)$, $v_z = \frac{1}{2}(v_x - iv_y)$). Using these notations, we can recast the coefficients in the following manner

$$\sigma_{xx} = \frac{2\pi e^2}{V} \text{Tr}[v_z \delta(E_F - H) v_{\bar{z}} \delta(E_F - H)] \quad (7)$$

$$\sigma_{xy}^I = \frac{ie^2}{\pi V} \text{Tr}[v_z G^- v_{\bar{z}} G^+ - v_z G^+ v_{\bar{z}} G^-] \quad (8)$$

$$\sigma_{xy}^{II} = \frac{ie^2}{2V} \text{Tr}[(zv_z - \bar{z}v_{\bar{z}}) \delta(E_F - H)] \quad (9)$$

In the expression of σ_{xy}^I , has been used the relation

$$\text{Tr} v_z G^\pm v_{\bar{z}} G^\pm = -\frac{1}{2} \text{Tr} G^\pm \quad (10)$$

a direct consequence of the commutation rules

$$v_z = \frac{i}{2} [\bar{z}, G^{-1}] \quad (11)$$

$$v_{\bar{z}} = \frac{i}{2} [z, G^{-1}] \quad (12)$$

2.2 Generating function

The Field theoretic approach to localization led to realize [29] the intimate relation between d.c. transport (at $T = 0$) and the symmetry between advanced and retarded Green function; the localization regime corresponds to the restauration by quantum interferences of this symmetry [30, 31], otherwise spontaneously broken by a non-vanishing density of states at the Fermi energy.

An explicit illustration of this, is the source-term formalism of Levine, Libby and Pruiskin [28], which expresses the d.c. transport coefficients as the response of the system to some rotation between the advanced and retarded wave. By convenience we will reformulate slightly this formalism in the following.

Consider the generating function

$$\mathcal{F}[\mathcal{A}_z^\alpha, \mathcal{A}_{\bar{z}}^\alpha] = \log \frac{\det[E_F - H(\mathcal{A}_z, \mathcal{A}_{\bar{z}}) + i\eta\tau_3]}{\det[E_F - H + i\eta\tau_3]} \quad (13)$$

It is left implicit here, that the operators involved in this expression acts on two-components wave functions, corresponding to either the retarded or advanced sector, as emphasized by the presence of the infinitesimal prescription $i\eta\tau_3$. The $\mathcal{A}_z^\alpha$ are the components of the sources $\mathcal{A}_z = \mathcal{A}_z^\alpha \tau_\alpha$, which are of matrix form, τ_α being the usual Pauli matrices

$$\tau_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \tau_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \tau_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (14)$$

In addition, we have

$$H(\mathcal{A}_z, \mathcal{A}_{\bar{z}}) = \left\{ \left(v_z - \frac{e}{2} \mathcal{A}_{\bar{z}} \right), \left(v_{\bar{z}} - \frac{e}{2} \mathcal{A}_z \right) \right\} + V(z, \bar{z}) \quad (15)$$

(where $\{, \}$ means anticommutator). If we restrict now the sources to be spatially uniform for $\alpha = 1, 2$ and

$$\mathcal{A}_z^3 = \frac{iB^3}{2} z \quad (16)$$

B^3 being a uniform magnetic field then, the static transport coefficients can be expressed in the following way

$$\sigma_{xx} = -\frac{1}{2\pi V} \frac{\partial^2 \mathcal{F}}{\partial \mathcal{A}_z^1 \partial \mathcal{A}_{\bar{z}}^1} \Big|_{\mathcal{A}=0} \quad (17)$$

$$\sigma_{xy}^I = -\frac{1}{\pi V} \frac{\partial^2 \mathcal{F}}{\partial \mathcal{A}_z^1 \partial \mathcal{A}_{\bar{z}}^2} \Big|_{\mathcal{A}=0} \quad (18)$$

$$\sigma_{xy}^{II} = -\frac{1}{2i\pi V} \frac{\partial \mathcal{F}}{\partial B^3} \Big|_{\mathcal{A}=0} \quad (19)$$

by direct comparison with (7,8,9), using the expansion

$$\mathcal{F}[\mathcal{A}_z^\alpha, \mathcal{A}_{\bar{z}}^\alpha] = \log \det[1 - \Delta H G] = -\text{Tr}[\Delta H G + \frac{1}{2}(\Delta H G)^2 + \dots] \quad (20)$$

with $\Delta H = H(\mathcal{A}_z, \mathcal{A}_{\bar{z}}) - H$ and $G = \begin{pmatrix} G^+ & 0 \\ 0 & G^- \end{pmatrix}$, and using the Ward identity (10) which in fact is a consequence of global $U(1) \times U(1)$ invariance of $\mathcal{F}$ [28]. The relation concerning σ_{xy}^{II} is reminiscent of the Streda formula (6). The connection is simply made by realizing that

$$\frac{\partial \mathcal{F}[\mathcal{A}_z^3, \mathcal{A}_{\bar{z}}^3]}{\partial E_F} = \text{Tr}[G^+(B + B^3) + G^-(B - B^3) - G^+(B) - G^-(B)] \quad (21)$$

(where the argument of the G 's denotes the sum of the magnetic field B and the fictitious one coming from the source $\mathcal{A}^3$). Since

$$G^\pm(B) = Re(B) \mp i\pi\rho(E_F, B) \quad (22)$$

($Re(B)$ is the real part of the Green function, and $\rho(E_F, B)$ is the density of states at the Fermi energy and for a given magnetic field B). Taking the first order term in the last relation, and integrating over energy leads to the relation

$$\frac{\partial \mathcal{F}}{\partial B^3} = -2i\pi V \frac{\partial N(E_F)}{\partial B} \quad (23)$$

which makes the connection with the Streda formula. The underlying reasons for these relations concerning σ_{xy}^{II} is that the source $\mathcal{A}^3$ is the conjugate field of the persistent current, locally given (at position $\mathbf{z} = (z, \bar{z})$) by

$$j_z^0(\mathbf{z}) = \frac{i}{2\pi} \text{Tr} [v_z | \mathbf{z} \rangle \langle \mathbf{z} | G \tau_3] \quad (24)$$

This entails, that if $\mathcal{A}^3$ corresponds to a magnetic field,

$$B^3 = \frac{1}{i} (\partial_z \mathcal{A}_z^3 - \partial_{\bar{z}} \mathcal{A}_{\bar{z}}^3) \quad (25)$$

then this magnetic field is conjugate to the magnetization (at the Fermi-energy), locally defined by

$$M(\mathbf{r}) = \frac{1}{2i\pi} \int d^2 \mathbf{z} \left(\frac{1}{\bar{z} - \bar{z}'} j_z^0(\mathbf{z}') - \frac{1}{z - z'} j_{\bar{z}}^0(\mathbf{z}') \right) \quad (26)$$

meaning that this quantity can be obtain from the generating function in the folowing manner

$$M(\mathbf{r}) = -\frac{1}{2i\pi} \frac{\partial \mathcal{F}}{\partial B^3(\mathbf{r})} \Big|_{A=0} \quad (27)$$

This is quite similar also to formulaes derived in the context of scattering theory [32, 33, 34] where the logarithm of the determinant of the scattering matrix is used as a generating function instead of $\mathcal{F}$.

3. The model

3.1 Pauli equation

After this preamble concerning the transport coefficients, we are now in position to define the model we are interested in, namely to specify equation (1). As already mentioned in the introduction, the random magnetic problem should be relevant for studying transport in a fractionnal quantum-Hall state, especially at even denominator filling factors. In this case, the ground-state consists of an assembly of composite Fermions, forming a weekly interacting Fermi-sea, and scattered by randomly distributed impurities. Due to the Chern-Simons gauge field needed to describe the ground-state, the effect of impurities is to produce fluctuations in the fictious magnetic field by modifying the local density of particules [1]. This entails the presence of magnetic vortices, located on the top of these impurities. For example, in a $\nu = \frac{1}{q}$ fractionnal state, a delta impurity is expected to be accompnied with q quantum of flux. As a consequence, a relevant model for this problem could actually be defined by

$$H = \frac{1}{2m} (\mathbf{p} - e\mathbf{A})^2 + \lambda B(\mathbf{r}) \quad (28)$$

where $B(\mathbf{r})$ is a random magnetic field plus the eventual effective external field, $\mathbf{A}$ the corresponding static gauge field, and λ a proportionality constant between the impurity scalar potential an the corresponding induced magnetic field. Once the statistical properties of the

magnetic field are specified, the model is entirely defined. Instead of studying this model for an arbitrary value of the parameter λ , we will restrict ourselves to the two particular values $\lambda = \pm \frac{e}{2m}$. In the context of the Aharonov-Bohm problem [35, 36], or Chern-Simons theory [37] these particular values correspond to the situation where the scale-invariance of the theory is not broken quantum mechanically and can be viewed as fixed point of the parameter λ under renormalization. They are moreover interpreted to be the spin-coupling constant of the electron with the magnetic field [38], the equation (28) beeing then precisely the Pauli equation, with the sign depending on the projection of the spin along the z axis. In the present context, this particular choice has as well the virtue of eliminating ultraviolet divergences, which otherwise occur at arbitrary λ in the perturbation expansion (of the average density of states for example), indicating also the scale invariance of the theory for these particular values. As a consequence we will consider this to be the most relevant situation, and postpone the analyses for arbitrary λ to further investigations.

For simplicity, we will consider a locally correlated random magnetic field, in its most simple expression, namely the white noise case,

$$\overline{B(\mathbf{r})B(\mathbf{r}')} = w\delta(\mathbf{r} - \mathbf{r}') \quad (29)$$

and with all the higher order cumulants set to zero. If we compare to a poissonian distribution of δ fluxes ϕ , with mean density n , the white noise distribution amounts to conserve only the second cumulant with a coefficient equal to

$$w = n\phi^2 \quad (30)$$

This approximation, to some extent, is justified from the field theoretical point of view, for scalar impurities (not magnetic). The second cumulant gives rise to a local quartic interaction term, and other cumulants contribute to terms of higher degree in the interaction, which are in principle irrelevant. This might be less justified in the random magnetic-field problem (see reference [8] for the average density of states for poissonian magnetic impurities, [9] for non-locally correlated poissonian magnetic impurities and [22] concerning the average conductivity for poissonian magnetic impurities), where as we shall see, the interacting term happens to be non-local.

3.2 Non-unitary transformation

The model beeing now defined let us consider it more explicitly. Given a magnetic field distribution $B(\mathbf{z})$, the static gauge field can be expressed in the Coulomb gauge ($\text{div}\mathbf{A} = 0$), using complex coordinates, according to

$$A_z = \frac{i}{2\pi} \int d^2\mathbf{z}' \frac{1}{\bar{z} - \bar{z}'} B(\mathbf{z}') \quad (31)$$

(recall that $\mathbf{z}$ means $(z, \bar{z})$, $B(\mathbf{z})$ is not supposed to be holomorphic). In the following we will have to distinguish between two situations, whether, in addition to the fluctuations, there is or not a uniform magnetic field.

When the average value of the magnetic field is non zero, then the two possible signs for the spin projection are unequivalent, and assuming by convention this uniform field to have the positive value B_0 , along the z axis, the Hamiltonian takes then the two possible forms

$$H^\pm = -2 \left(\partial_z - \frac{1}{4\pi} \int d^2\mathbf{z}' \frac{1}{z - z'} B(\mathbf{z}') \right) \left(\partial_{\bar{z}} + \frac{1}{4\pi} \int d^2\mathbf{z}' \frac{1}{\bar{z} - \bar{z}'} B(\mathbf{z}') \right) + (1 \pm 1) B(\mathbf{z}') \quad (32)$$

Where the value of e and m are from now on set to 1. Having in mind to average later on over disorder, we perform a non-unitary transformation on the wave-function, in order to eliminate the quadratic term in the fluctuating magnetic field [8]

$$\psi(\mathbf{z}) = e^{\pm \frac{1}{2\pi} \int d^2\mathbf{z} b(\mathbf{z}) \log |z-z'|} \tilde{\psi}(\mathbf{z}) \quad (33)$$

$b = (B - B_0)/B_0$ is the dimensionless fluctuating magnetic field. The generating function $\mathcal{F}$ which will be used later, is invariant under this transformation, as long as the Hilbert space is not modified. This is indeed the case, qualitatively because $b(\mathbf{z})$ average itself to zero over the plane, consequently wave functions remain square integrable under this transformation, or under the inverse. At this point it is convenient to introduce the raising and lowering Landau level operators,

$$a = \partial_z + \frac{1}{2}z \quad (34)$$

$$a^\dagger = -\partial_{\bar{z}} + \frac{1}{2}\bar{z} \quad (35)$$

(the cyclotron energy is set to one) The Hamiltonian $\tilde{H}^\pm$ acting on the wave function $\tilde{\psi}$ takes then the simple form

$$\tilde{H}^+ = 2aa^\dagger + \frac{1}{\pi} \int d^2\mathbf{z}' \frac{b(\mathbf{z}')}{\bar{z} - \bar{z}'} a^\dagger \quad (36)$$

$$\tilde{H}^- = 2a^\dagger a - \frac{1}{\pi} \int d^2\mathbf{z}' \frac{b(\mathbf{z}')}{z - z'} a \quad (37)$$

This last expression, corresponding to the spin down case, makes the (expected) zero mode directly apparent. It consists of the lowest Landau level itself, giving therefore a degeneracy to these particular states in accordance with the Atiyah-Singer theorem. The fact that this feature explicitly appears here, is due to the very analogous form of the factor multiplying $\tilde{\psi}$ with the Aharonov-Casher solution [39].

In absence of a uniform magnetic field, considering the statistical properties of the random field, there is no privileged orientation given to the plane. The sign of the spin projection is therefore indifferent, and taking it negative by convention, we then perform the same non-unitary transformation (minus sign in the exponent). The Hamiltonian acting on $\tilde{\psi}$ has then the expression

$$\tilde{H} = -2\partial_z \partial_{\bar{z}} + \frac{1}{\pi} \int d^2\mathbf{z}' \frac{b(\mathbf{z}')}{\bar{z} - \bar{z}'} \partial_z \quad (38)$$

4. Saddle point DOS

4.1 Replica Supersymmetry representation

We turn now to the functional approach to the question of computing the d.c. transport coefficients. The objective is to average the generating function over disorder, which necessitates to express it in a convenient form. Although we know that in principle, in order to compute the mean conductivity, it is sufficient to average $\exp(\mathcal{F})$ (the equivalent of the partition function, instead of the free energy), we will follow the logic of this presentation, and average $\mathcal{F}$ itself. Using the combination of the supersymmetry method of Efetov [31] and the replica trick, it is possible to express $\mathcal{F}$ in a suitable form for average.

The supersymmetric calculus gives the possibility to view a ratio of two determinants as a superdeterminant

$$\frac{\det[E_F - H(\mathcal{A}_z, \mathcal{A}_{\bar{z}}) + i\eta\tau_3]}{\det[E_F - H + i\eta\tau_3]} = S \det[E - H(\mathcal{A}_z, \mathcal{A}_{\bar{z}}) + i\eta\tau_3] \quad (39)$$

where the numerator corresponds to the grassmann sector and the denominator to the bozonic one. The sources are still ordinary complex fields, but are acting in the Grassmann sector. The superdeterminant is then expressed in terms of a Gaussian integral over superfields. Moreover, the replica-trick amounts to express the logarithm according to

$$\log S \det[E - H(\mathcal{A}_z, \mathcal{A}_{\bar{z}}) + i\eta\tau_3] = \lim_{q \rightarrow 0} \frac{d}{dq} S \det[E - H(\mathcal{A}_z, \mathcal{A}_{\bar{z}}) + i\eta\tau_3]^q \quad (40)$$

which has to be understood by means of analytical continuation of q to real values. This allows to express $\mathcal{F}$ in the way

$$\mathcal{F}[\mathcal{A}_z^\alpha, \mathcal{A}_{\bar{z}}^\alpha] = \lim_{q \rightarrow 0} \frac{d}{dq} \int D\bar{\Psi} D\Psi e^{i \int d^2\mathbf{z} \bar{\Psi}(E - H(\mathcal{A}_z, \mathcal{A}_{\bar{z}}) + i\eta\tau_3 \times I_q) \Psi} \quad (41)$$

where the integration is performed over $2q$ components superfields, q complexes and q grassman components. In this form it is now possible to average over the magnetic fluctuations. Let us consider the case with no uniform magnetic field. The term to be averaged is

$$\langle e^{-\frac{i}{4\pi} \int d^2\mathbf{z} d^2\mathbf{z}' \bar{\Psi}(\mathbf{z}) \frac{b(\mathbf{z}')}{\bar{z} - \bar{z}'} D_z \Psi(\mathbf{z})} \rangle_b = e^{\frac{w}{2\pi} \int d^2\mathbf{z}_1 d^2\mathbf{z}_2 \bar{\Psi}(\mathbf{z}_1) D_{z_1} \Psi(\mathbf{z}_1) \frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} \bar{\Psi}(\mathbf{z}_2) D_{z_2} \Psi(\mathbf{z}_2)} \quad (42)$$

where D_z replaces ∂_z because of the sources and where has been utilized the identity

$$\int d^2\mathbf{z}' \frac{1}{(\bar{z}_1 - \bar{z}')(\bar{z}_2 - \bar{z}')} = -\pi \frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} \quad (43)$$

As already anticipated, the field theory we are obtaining, contains, in addition to the free part of the action,

$$\mathcal{S}_0 = i \int d^2\mathbf{z} \bar{\Psi}(\mathbf{z}) (E + 2D_z D_{\bar{z}} + i\eta\tau_3 \times I_q) \Psi(\mathbf{z}) \quad (44)$$

a quartic interacting term, which in contrary to the usual random scalar potential, happens to be non local, because of the long range nature of the Aharonov-Bohm interaction. This term can be reexpressed using the definition of the supertrace

$$\mathcal{S}_{int} = \frac{w}{2\pi} \int d^2\mathbf{z}_1 d^2\mathbf{z}_2 Str \left[\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} D_{z_1} \Psi(\mathbf{z}_1) \bar{\Psi}(\mathbf{z}_2) D_{z_2} \Psi(\mathbf{z}_2) \bar{\Psi}(\mathbf{z}_1) \right] \quad (45)$$

and can nevertheless be decoupled, using a Hubbard-Strattanovich transformation, that is to say the identity

$$e^{\mathcal{S}_{int}} = \int D\bar{Q} DQ e^{-\frac{2\pi}{w} Str \bar{Q} Q + \int d^2\mathbf{z}_1 d^2\mathbf{z}_2 Str \left[\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} D_{z_1} \Psi(\mathbf{z}_1) \bar{\Psi}(\mathbf{z}_2) Q(\mathbf{z}_2, \mathbf{z}_1) + \bar{Q}(\mathbf{z}_1, \mathbf{z}_2) D_{z_2} \Psi(\mathbf{z}_2) \bar{\Psi}(\mathbf{z}_1) \right]} \quad (46)$$

The functionnal integration involve $2n \times 2n$ non-local supermatrices with no particular constraint, in contrast to the local theory [31]. Permutating again the supertrace in $\mathcal{S}_{int}$, and integrating over the $2n$ components superfields, we obtain

$$\mathcal{F}[\mathcal{A}_z^\alpha, \mathcal{A}_{\bar{z}}^\alpha] = \lim_{q \rightarrow 0} \frac{d}{dq} \int D\bar{Q} DQ e^{Str \log \left[E + 2D_z D_{\bar{z}} + i\eta\tau_3 \times I_q + i \frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} Q D_z + i \bar{Q} D_{\bar{z}} \right] - \frac{2\pi}{w} Str \bar{Q} Q} \quad (47)$$

4.2 Saddle point approximation

The saddle point is then obtained as the stationnary point when Q and $\bar{Q}$ are varied independantly. Expanding at first order the supertrace of the logarithm in the above expression, leads to the set of equations

$$Q_0(\mathbf{z}_1, \mathbf{z}_2) = i \frac{w}{2\pi} \langle \mathbf{z}_1 | \partial_z G_0 | \mathbf{z}_2 \rangle \quad (48)$$

$$\bar{Q}_0(\mathbf{z}_1, \mathbf{z}_2) = i \frac{w}{2\pi} \frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} \langle \mathbf{z}_1 | \partial_z G_0 | \mathbf{z}_2 \rangle \quad (49)$$

$$G_0(\mathbf{z}_1, \mathbf{z}_2) = \langle \mathbf{z}_1 | \frac{1}{E + 2\partial_z \partial_{\bar{z}} + 2i\bar{Q}_0 \partial_z} | \mathbf{z}_2 \rangle \quad (50)$$

the sources beeing set to zero for the moment. The main difference with the scalar case is that these relations are equalities between operators. In absence of a uniform magnetic field, we are seeking a solution which preserves the translationnal and rotationnal invariance, meaning that G has to be a function of $p = |\mathbf{p}|$. In addition we look for this solution in a diagonal form. In that case the equation for any diagonal components $g(E, p)$ of $G(\mathbf{p})$ rewrites

$$g(E, p) = \frac{1}{E - \frac{1}{2}p^2 - \frac{w}{4\pi}p^2 g(E, p)} \quad (51)$$

This equation admits two real solutions

$$g^\pm(E, p) = \frac{2\pi}{wp^2} \left(E - \frac{1}{2}p^2 \pm \sqrt{(E - E_p^-)(E - E_p^+)} \right) \quad (52)$$

outside the energy interval $[E_p^-, E_p^+]$ with

$$E_p^\pm = \frac{1}{2}p^2 \pm \sqrt{\frac{w}{\pi}}p \quad (53)$$

One corresponds to a stable saddle point, the other to an unstable one. By looking at the sequence $g_{n+1}(E, p) = 1/(E - \frac{1}{2}p^2 - \frac{w}{4\pi}p^2 g_n(E, p))$ one observe that when $E < E_p^-$, the attractive fixe point (stable) is $g^+(E, p)$, even though it is $g^-(E, p)$ when $E > E_p^+$. This is then sufficient to analytically continue $g(E, p)$ onto the complexe plane, without any ambiguity,

$$g(Z, p) = \frac{2\pi}{wp^2} \left(Z - \frac{1}{2}p^2 - \sqrt{(Z - E_p^-)}\sqrt{(Z - E_p^+)} \right) \quad (54)$$

where the square root is defined by using the convention that the argument of $Z - E_p^-$ and $Z - E_p^+$ vary in the interval $[-\pi, \pi]$. As a result we have a branch cut situated on the segment $Z \in [E_p^-, E_p^+]$,

$$g(E \pm i\eta, p) = \frac{2\pi}{wp^2} \left(E - \frac{1}{2}p^2 \mp i\sqrt{(E_p^+ - E)(E - E_p^-)} \right) \quad (55)$$

corresponding to a non-vanishing contribution to the density of states

$$\rho_p(E) = \frac{1}{2i\pi} (G_p(E - i\eta) - G_p(E + i\eta)) = \frac{2}{wp^2} \sqrt{(E_p^+ - E)(E - E_p^-)} \quad (56)$$

Therefore, the SCBA density of states is obtained by summing this last expression with the measure $pd\mathbf{p}/2\pi$, which leads to

$$\rho(E) = \begin{cases} 0 & E < -\frac{w}{2\pi} \\ \frac{1}{2w}(2E + \frac{w}{\pi}) & -\frac{w}{2\pi} < E < 0 \\ \frac{1}{2\pi} & E > 0 \end{cases} \quad (57)$$

and is responsible for the spontaneous breaking of the symmetry $V(2n) \rightarrow V(n) \times V(n)$, i.e. between the advanced and retarded sector, at the saddle-point

$$G_0(E, p) = \frac{2\pi}{wp^2} \left(E - \frac{1}{2}p^2 - i\tau_3 \times I_q \sqrt{(E_p^+ - E)(E - E_p^-)} \right) \quad (58)$$

The negative part of the DOS is actually unphysical, because the spectrum of this problem is supposed to be positive. The possible origin of this undesirable feature shall be discussed in the conclusive section.

4.3 Broadened Landau spectrum

If we turn now to the case, where a uniform magnetic field is present, we are expecting to find a saddle-point solution which is both invariant under magnetic translation and rotations around the origin (if we choose the symmetric gauge centered at the origin). The representation of this group of transformation is irreducible in each Landau level, meaning that $G(E)$ has to be decomposed on the set of projection operator P_n corresponding to each Landau Level. In other words, $G(E)$ has to be a function of $aa^\dagger$

$$G(E, aa^\dagger) = \sum_{n=1}^{\infty} G_n(E) P_n \quad (59)$$

For the spin-up case, the saddle point equation (with the sources set to zero again) takes the form

$$G(\mathbf{z}_1, \mathbf{z}_2) = \langle \mathbf{z}_1 | \frac{1}{E - 2aa^\dagger - \frac{w}{\pi} \frac{z-z'}{\bar{z}-\bar{z}'} a^\dagger} G a^\dagger | \mathbf{z}_2 \rangle \quad (60)$$

(where a shorthand notation has been to express that the kernel of $a^\dagger G a^\dagger$ is multiplied by $\frac{z-z'}{\bar{z}-\bar{z}'}$). Using the properties (see appendix) that

$$\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} \langle \mathbf{z}_1 | a^\dagger G(E, aa^\dagger) a^\dagger | \mathbf{z}_2 \rangle = \langle \mathbf{z}_1 | aa^\dagger G(E, aa^\dagger) | \mathbf{z}_2 \rangle \quad (61)$$

and following the lines of reasoning described in the preceeding case, we obtain (with n labelling the set of Landau levels)

$$G_0(E, n) = \pi^2 (e_n(E) - i\tau_3 \times I_q \rho_n(E)) = \frac{2}{E - 2n + i\tau_3 \times I_q \sqrt{(E - E_n^-)(E_n^+ - E)}} \quad (62)$$

with

$$E_n^\pm = 2n \pm 2\sqrt{\frac{w}{\pi}n} \quad (63)$$

which leads to the standard semi-circle law for the SCBA density of states

$$\rho_n(E) = \frac{1}{2\pi^2 n \alpha} \begin{cases} \sqrt{4\alpha n - (E/b - 2n)^2} & E_n^- < E < E_n^+ \\ 0 & \text{otherwise} \end{cases} \quad (64)$$

where the parameter $\alpha = \frac{w}{b\pi}$ has been introduced, and where the dependency with the b field has been restored. In this form it is explicitly seen that $\rho(E)$, and actually any other quantity is a function of E/b and α , the scaling property of the system is not broken by any UV cut-off. The real part of the Green function is given by

$$e_n(E) = \frac{1}{2\pi^2 n \alpha} \begin{cases} E/b - 2n + \sqrt{(E/b - 2n)^2 - 4n\alpha} & E < E_n^- \\ E/b - 2n & E_n^- < E < E_n^+ \\ E/b - 2n - \sqrt{(E/b - 2n)^2 - 4n\alpha} & E > E_n^+ \end{cases} \quad (65)$$

The DOS is therefore composed of a sequence of enlarged Landau levels, with a width increasing in proportion to the square root of the LL index n . The result is not very different from what is obtained for a poissonian magnetic impurities system [8, 10] This is however different from the scalar random potential situation, where the width is not n dependant, and this fact has a simple semi-classic interpretation [40, 9]; the semi-classic orbits in the Landau problem consists of cyclotron orbits, the number of time the electron circulates around such an orbite corresponds to which Landau level it belongs to. Moreover, the magnetic flux passing through a given area,




Figure 1: Broadening of the Landau Levels at fixed b for the values $\alpha = 10^{-2}, 310^{-2}, 10^{-1}, 1, 10$ of the disorder parameter.

has fluctuations proportionnal to the square root of this area, meaning that the fluctuations of the flux passing through a cyclotron orbit surrounded n times, is proportionnal to $\sqrt{n}$, giving the typical width of the broadened Landau levels. In the spin-down case, the saddle point equation reads

$$G(\mathbf{z}_1, \mathbf{z}_2) = \langle \mathbf{z}_1 | \frac{1}{E - 2a^\dagger a - \frac{w}{\pi} \frac{\bar{z} - \bar{z}'}{z - z'} a} G a | \mathbf{z}_2 \rangle \quad (66)$$

and using now the property

$$\frac{\bar{z}_1 - \bar{z}_2}{z_1 - z_2} \langle \mathbf{z}_1 | a G(E, aa^\dagger) a | \mathbf{z}_2 \rangle = \langle \mathbf{z}_1 | a^\dagger a G(E, aa^\dagger) | \mathbf{z}_2 \rangle \quad (67)$$

we obtain the same expressions but with the modification $n \rightarrow n - 1$. In particular, the lowest Landau level at $E = 0$ remains unperturbed.

5. Saddle point Conductivity

5.1 In absence of a uniform magnetic field

The next step is to calculate the transport coefficient at the saddle point. Using the generating function, it amounts to replace the integration over the Q and $\bar{Q}$ matrices in (47) by the saddle-point solution Q_0 and $\bar{Q}_0$ of equation (48). Since the saddle point doesn't break the replica symmetry, we therefore obtain the saddle point expression for the generating function

$$\mathcal{F}_0[\mathcal{A}_z^\alpha, \mathcal{A}_{\bar{z}}^\alpha] = \log S \det \left[E + 2D_z D_{\bar{z}} + 2i\bar{Q}_0 D_z \right] \quad (68)$$

using the fact that $\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} Q_0(\mathbf{z}_1, \mathbf{z}_2) = \bar{Q}_0(\mathbf{z}_1, \mathbf{z}_2)$. By differentiation with respect to the sources, according to recipes given in part one, we would then obtain bare values for transport coefficients. However, at this point, we have to realize that the saddle point generating function doesn't fulfil local $U(1) \times U(1)$ (gauge) invariance, namely because Q_0 and $\bar{Q}_0$ are non-local operators. This originates in their dependancy with respect to the operators ∂_z and $\partial_{\bar{z}}$ accordingly to equation (48). A straightforward procedure to restore gauge invariance would be then, to replace these operators by D_z and $D_{\bar{z}}$. However, this procedure is ambiguous because τ_3 does not commute with D_z and $D_{\bar{z}}$, leaving any expression of Q_0 to be arbitrary. Instead, if we compute the saddle point, while keeping the sources fixed, we obtain the correct dependancy of Q_0 and $\bar{Q}_0$ with the sources. Let us denote $Q(\mathcal{A})$ and $\bar{Q}(\mathcal{A})$ the corresponding saddle point solutions. From the first part we know that it is sufficient to consider uniform sources to compute the conductivity tensor. In that case the saddle point expression for the generating function can be written in the form

$$\mathcal{F}_0[\mathcal{A}_z^\alpha, \mathcal{A}_{\bar{z}}^\alpha] = \log S \det \left[E + \{D_z, D_{\bar{z}}\} + i\bar{Q}(\mathcal{A})D_z - iD_{\bar{z}}Q(\mathcal{A}) \right] - \frac{2\pi}{w} \text{Str} \bar{Q}(\mathcal{A})Q(\mathcal{A}) \quad (69)$$

with the saddle point constraint

$$Q(\mathcal{A}) = i\frac{w}{2\pi} D_z G(\mathcal{A}) \quad (70)$$

$$\bar{Q}(\mathcal{A}) = -i\frac{w}{2\pi} G(\mathcal{A})D_{\bar{z}} \quad (71)$$

$$G(\mathcal{A}) = \frac{1}{E + \{D_z, D_{\bar{z}}\} + \frac{w}{2\pi} \{G(\mathcal{A}), D_{\bar{z}}D_z\}} \quad (72)$$

The computation of transport coefficients necessitates then to solve this last equation. Assuming that $G(\mathcal{A})$ is analytic with respect to the sources, it is then sufficient to solve at first order in the source. Consider the expansion

$$G(\mathcal{A}) = G_0 + G_\alpha^z \mathcal{A}_z^\alpha + G_\alpha^{\bar{z}} \mathcal{A}_{\bar{z}}^\alpha + .. \quad (73)$$

Deriving (69) twice with respect to the sources, according to the saddle point constraints leads to the following expression for the conductivity coefficients

$$\sigma_{xx} = \frac{1}{4\pi V} \text{Tr} \left\{ G_0 + \frac{w}{4\pi} G_0^2 - 2 \left[v_{\bar{z}} + \frac{w}{4\pi} (v_{\bar{z}} G_0 + G_0^\dagger v_{\bar{z}}) \right] G_1^z \tau_1 \right\} \quad (74)$$

$$\sigma_{xy}^I = \frac{i}{2\pi V} \text{Tr} \left\{ \frac{w}{4\pi} G_0^2 - 2 \left[v_{\bar{z}} + \frac{w}{4\pi} (v_{\bar{z}} G_0 + G_0^\dagger v_{\bar{z}}) \right] G_1^z \tau_1 \right\} \tau_3 \quad (75)$$

where the trace is yet restricted to the Grassmann sector, and $G_0^\dagger$ on this sector reads

$$G_0^\dagger = \begin{pmatrix} G_0^- & 0 \\ 0 & G_0^+ \end{pmatrix} \quad (76)$$

whereas

$$G_0 = \begin{pmatrix} G_0^+ & 0 \\ 0 & G_0^- \end{pmatrix} \quad (77)$$

In absence of a net magnetic field, we expect G_α^z to be a function of p and $\bar{p}$ because of translation invariance. Solving (72) with this assumption we obtain

$$G_\alpha^z(\mathbf{p}) = -\frac{\bar{p}}{2} \frac{\left(1 + \frac{w}{4\pi}(G_0 + G_0^\dagger)\right) G_0 G_0^\dagger}{1 - \frac{w}{4\pi} p^2 G_0 G_0^\dagger} \tau_\alpha \quad \alpha = 1, 2 \quad (78)$$

the striking feature of this last expression is that it is not defined when p is such that $E_p^- < E_F < E_p^+$, the denominator vanishes because in this interval we have (55)

$$G_0^+ G_0^- = \frac{4\pi}{w p^2} \quad (79)$$

The reason for this is that the procedure of keeping sources in the computation of the saddle point, amounts to incorporate ladder diagrams contribution, when the trace is taken. These diagrams are in principle responsible for weak localization corrections, and since the system, because of the magnetic fluctuations, is not time-reversal invariant, they are actually the only ones. Therefore this divergency indicates that $G(\mathcal{A})$ is not an analytic function with respect to the sources, which has to be related to the eventuality that the system is localized. The expression of $G_\alpha^z(\mathbf{p})$ suggests the possibility of a self-consistent treatment of the problem [41]; this would however require a more complete treatment, beyond saddle point approximation, which is out of the scope of this paper.

5.2 Conductivity in presence of a uniform magnetic field

The situation is quite different when an external uniform magnetic field is added to the problem. In that case, the IR divergency observed preceedingly happens to be removed. Indeed, the solution to (72) can be searched in the form

$$G_\alpha^z(a, a^\dagger) = a^\dagger g(aa^\dagger) \tau_\alpha \quad (80)$$

and using the rules (for an arbitrary function f),

$$a f(a^\dagger a) = f(aa^\dagger) a \quad (81)$$

$$a^\dagger f(aa^\dagger) = f(a^\dagger a) a^\dagger \quad (82)$$

we obtain the solution

$$G_\alpha^z(a, a^\dagger) = -i \frac{\left[1 + \frac{w}{4\pi} (G_0(aa^\dagger) + G_0^\dagger(a^\dagger a))\right] G_0(aa^\dagger) G_0^\dagger(a^\dagger a)}{1 - \frac{w}{2\pi} (2aa^\dagger - 1) G_0(aa^\dagger) G_0^\dagger(a^\dagger a)} a^\dagger \tau_\alpha \quad (83)$$

for $\alpha = 1, 2$ and

$$G_3^z(a, a^\dagger) = -i \frac{\left[1 + \frac{w}{4\pi} (G_0(aa^\dagger) + G_0(a^\dagger a))\right] G_0(aa^\dagger) G_0(a^\dagger a)}{1 - \frac{w}{2\pi} (2aa^\dagger - 1) G_0(aa^\dagger) G_0(a^\dagger a)} a^\dagger \tau_3 \quad (84)$$




Figure 2: Transverse conductivity in unit of $\frac{e^2}{h}$ for the spin-up (left) and spin-down (right) cases for $\alpha = 10^{-2}, 10^{-1}, 1, 10$, compared to the classical value (straight line). The x axes is the Landau level filling number.

where $G_0(aa^\dagger)$ is defined according to (59) and (62). Inserting this last expression in (87) and (89) allows to compute the conductivity tensor. However for σ_{xx} it is convenient to use the following Ward identity which is a consequence of global $U(1) \times U(1)$ invariance,

$$Tr \left\{ G_0 + \frac{w}{4\pi} G_0^2 - 2 \left[v_{\bar{z}} + \frac{w}{4\pi} (v_{\bar{z}} G_0 + G_0 v_{\bar{z}}) \right] G_3^z \tau_3 \right\} = 0 \quad (85)$$

and which is the saddle point version of (10). Defining

$$g_n^s = \pi^2 (e_n(E_F) - i s \rho_n(E_F)) \quad (86)$$

with $s = \pm$ and $e_n(E_F)$ and $\rho_n(E_F)$ given by (65) and (64) we can then express σ_{xx} in the form (recall $\alpha = \frac{w}{b\pi}$)

$$\sigma_{xx} = -\frac{1}{2\pi^2} \sum_{s,s'} \sum_{n=1}^{\infty} s s' (n-1) \frac{\left(1 + \frac{\alpha}{4} (g_n^s + g_{n-1}^{s'})\right)^2 g_n^s g_{n-1}^{s'}}{1 - \frac{\alpha}{2} (2n-1) g_n^s g_{n-1}^{s'}} \quad (87)$$

In the strong magnetic field limit, i.e. when the Landau Levels do not overlap, it is quite easy to see that σ_{xx} vanishes. Surprisingly enough this happens to be true whatever b is (numerical check)

$$\sigma_{xx} = 0 \quad (88)$$

Concerning the transverse conductivity, we obtain for the dissipative part

$$\sigma_{xy}^I = \sum_{n=1}^{\infty} \left\{ \frac{1}{4n} (E_F/b - 2n) \rho_n(E_F) + \frac{i}{\pi^2} \sum_{s=\pm} s (n-1) \frac{\left(1 + \frac{\alpha}{4} (g_n^s + g_{n-1}^{-s})\right)^2 g_n^s g_{n-1}^{-s}}{1 - \frac{\alpha}{2} (2n-1) g_n^s g_{n-1}^{-s}} \right\} \quad (89)$$

although the contribution coming from the magnetization is obtained by using the Streda formula (6)

$$\sigma_{xy}^{II} = \frac{1}{b} N(E_F) - \sum_{n=1}^{\infty} \frac{1}{2} (E_F/b + 2n) \rho_n(E_F) \quad (90)$$

Altogether the sum simplifies exactly to

$$\sigma_{xy}^{up} = \frac{1}{b}N(E_F) - \sum_{n=1}^{\infty} \rho_n(E_F) = \frac{1}{b}N(E_F) - \rho(E_F) \quad (91)$$

For the spin-down case, all the n appearing in the fraction of expression (87) and (89) are shifted by -1 . The conclusion concerning σ_{xx} is not modified, the longitudinal conductivity vanishes exactly for all values of the magnetic field. For σ_{xy} we obtain a slightly modified expression, except for the sign relative to the “classical” value,

$$\sigma_{xy}^{down} = \frac{1}{b}N(E_F) + \sum_{n=1}^{\infty} \frac{E_F}{2bn} \rho_n(E_F) \quad (92)$$

which again is obtained with no approximation. The Hall conductance is plotted in figure 2 for the two orientations of the spin. In the strong magnetic field limit we obtain oscillations of the transverse conductivity, in qualitative agreement with the conclusion of reference [22] for the spin-down case. The spin-up plot resembles qualitatively the numerical results obtained in reference [42].




Figure 3: Difference between the spin-up and the spin-down transverse conductivity in unit of $\frac{e^2}{h}$ for $b = 5, 1, .1, .01$ and $w = \pi$, versus Fermi energy.

The limit of vanishing mean magnetic field is a bit more mysterious. It should be first noticed that σ_{xy} as a function of electronic integrated density is divergent. Although paradoxical, one can consider this to be coherent with the fact that σ_{xx} vanishes, if one think of the semi-classical relation between these two coefficients [43]. Anyway, something interesting happens when we compare the Hall conductance for the two different spin projections in this limit, as a function of the Fermi energy (see figure 3). This difference takes the quantized value of 2 (in unit of $\frac{e^2}{h}$) when b goes to zero. The first quantum is simply traced back to the additionnal Landau level lying at $E = 0$ in the spin-down situation and which is not affected by disorder (see the strong magnetic field curve $b = 5$ in figure 3). The second one seems to indicate that the system restores the chiral symetry (considering spin-up and spin down situations as a whole) when the average b goes to zero, by reintroducing progressively a zero energy state in the spin-up

spectrum. In addition this state restores parity, by contributing with the opposite sign to the Hall conductance. In this sense, the limit of zero magnetic field is unstable with respect to spin-flip at $E = 0$.

6. Conclusion

The formalism presented in this paper, which proposes to compute directly static transport coefficients from a generating function, has been used at the basic level of the saddle point approximation. Already at this stage was exemplified the possibility of incorporating first dominant contributions, namely the ladder diagrams contribution, by keeping uniform sources when computing the saddle point. In the example we considered, the white noise random magnetic field, it was observed that in absence of an external uniform magnetic field, the saddle point solution is not analytic with respect to the sources indicating a non-simply diffusive behavior. Slowly spatially varying sources should be considered in order to analyse further this question, and a self-consistent argument analogous to the one of Vollhardt and Wölfle for the scalar disorder case [41] could be elaborated. Furthermore, this formalism should be possibly generalized for the computation of dynamical transport coefficients, and eventually extended to the non-linear regime.

Concerning the random magnetic field problem, the free parameter λ which was introduced in section 3, was only considered for two particular values, which presumably insures the system to remain scale invariant quantum mechanically. The fact that the longitudinal conductivity was suprizingly found to remain exactly zero, when Landau levels do overlap, although not really conclusive (within the saddle-point approximation), is in favour of this hypothesis. Interestingly enough, although may be more technically involved, would be to consider this model for arbitrary λ , in particular to see wether it is possible to find at this level of approximation a relation between σ_{xx} and σ_{xy} analogous to the one obtained by Ando et al. for the scalar random potential [43]. In addition a flow between these two points under scale transformation might be expected. Beforehand the problem of undesirable negative energy states which appear in zero magnetic field has to be cured, may be by avoiding the non-unitary transformation, which presumably causes some non-normalizable states to re-enter the Hilbert space. This could possibly clarify the nature of the zero energy state in the average magnetic field limit, which, as suggested by the analyses of figure 3 seems to spontaneously break parity and to be able to carry one quantum of Hall current.

Acknowledgements

I am pleased to thank J. Desbois, S. Ouvry and C. Texier for useful discussions concerning their approach to this problem, J.M. Leinaas, and H.A. Weidenmueller for discussing the supersymmetric approach to the transport problem. I am thankful to the Institute of physics of the Oslo University for hospitality, and the National Reasearch Council of Norway for financial support.

Appendix

In this section we justify the use of the formula

$$\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} \langle \mathbf{z}_1 | a^\dagger f(aa^\dagger) | \mathbf{z}_2 \rangle = \langle \mathbf{z}_1 | f(aa^\dagger) a | \mathbf{z}_2 \rangle \quad (93)$$

or equivalently

$$\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} \langle \mathbf{z}_1 | f(aa^\dagger) a^\dagger | \mathbf{z}_2 \rangle = \langle \mathbf{z}_1 | af(aa^\dagger) | \mathbf{z}_2 \rangle \quad (94)$$

where f is an arbitrary function of $aa^\dagger$, which is actually defined by the values f_n it takes on each Landau Level

$$f(aa^\dagger) = \sum_{n=1}^{\infty} f_n P_n \quad (95)$$

To this end, it is convenient to introduce the raising and lowering operators of angular momentum

$$b = \partial_z + \frac{1}{2}\bar{z} \quad (96)$$

$$b^\dagger = -\partial_{\bar{z}} + \frac{1}{2}z \quad (97)$$

and a generating function for the set of projection operators on the Landau Levels, namely the thermal propagator

$$G_t = e^{-aa^\dagger t} = \sum_{n=1}^{\infty} e^{-nt} P_n \quad (98)$$

Then, according to the commutation rules between the a , $a^\dagger$ and b , $b^\dagger$ operators we have the property

$$[\hat{z}, G_t] = [a + b^\dagger, G_t] = [a, G_t] = (1 - e^t)aG_t \quad (99)$$

as a consequence, it turns out that

$$\langle \mathbf{z}_1 | aG_t | \mathbf{z}_2 \rangle = \frac{e^{-t}}{1 - e^{-t}} (z_1 - z_2) \langle \mathbf{z}_1 | G_t | \mathbf{z}_2 \rangle \quad (100)$$

In the same way we have

$$\langle \mathbf{z}_1 | a^\dagger G_t | \mathbf{z}_2 \rangle = \frac{1}{1 - e^{-t}} (\bar{z}_1 - \bar{z}_2) \langle \mathbf{z}_1 | G_t | \mathbf{z}_2 \rangle \quad (101)$$

which leads to obtain

$$\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} \langle \mathbf{z}_1 | a^\dagger G_t | \mathbf{z}_2 \rangle = \langle \mathbf{z}_1 | G_t a | \mathbf{z}_2 \rangle \quad (102)$$

or equivalently

$$\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2} \langle \mathbf{z}_1 | G_t a^\dagger | \mathbf{z}_2 \rangle = \langle \mathbf{z}_1 | a G_t | \mathbf{z}_2 \rangle \quad (103)$$

G_t being a generating function for the Landau Levels projection operator, the formula is therefore valid for an arbitrary function of $aa^\dagger$ as stated above.

References

- [1] B.I. Halperin, P.A. Lee and N. Read, Phys. Rev. B **47** 7312;
- [2] L.B. Ioffe and V. Kalmeyer, Phys. Rev. B **44** (1991) 750;
- [3] B.L. Altshuler, L.B. Ioffe, Phys. Rev. Lett. **69** (1992) 2979;
- [4] G. Gavazzi, J.M. Wheatley, and A.J. Shofield, Phys. Rev. B **47** (1993) 15170;

- [5] D. V. Khveshenko and S.V. Meshkov, Phys. Rev B **47** (1993) 12051;
- [6] A. Lusakowski and L.A. Turski Phys. Rev. B **48** (1993) 3835;
- [7] A. Barelli, R. Fleckinger and T. Ziman, Phys. Rev. B **49** (1994) 3340; Europhys. Lett. **27** (1994) 531;
- [8] J. Desbois, C. Furtlehner and S. Ouvry, Nucl. Phys B **453**[FS] (1995) 759; J. Phys. I France **6** (1996) 641;
- [9] J. Desbois, C. Furtlehner and S. Ouvry, J. Phys. A: Math. Gen. **30** (1997) 7291;
- [10] C. Furtlehner, Thèse de 3^{ème} cycle, Université Paris 6
- [11] J. Rammer and A.L. Shelankov, Phys. Rev. B **36**(1987) 3135;
- [12] A.G. Aronov, A.D. Mirlin, P. Wölfle, Phys. Rev. B **49**(1994) 16609;
- [13] S.C. Zhang and D.P. Arovas, Phys. Rev. Lett. **72** (1994) 1886;
- [14] C. Pryor and A. Zee, Phys. Rev. B **46** (1992) 3116;
- [15] Y. Avishai, Y. Hatsugai and M. Kohmoto, Phys. Rev. B **47** (1993) 9561;
- [16] D.Z. Liu, X.C. Xie, S. Das Sarma and S.C. Zhang, Phys. Rev. B **52** (1995) 5858;
- [17] J. Miller and J. Wang, Phys. Rev. Lett. **76** (1996) 1461;
- [18] A. Furusaki, Phys. Rev. Lett. **82** (1999) 604;
- [19] A. Altland and B.D. Simons, preprint cond-mat/9811134 (1999);
- [20] A. Lusakowski and L.A. Turski, preprint cond-mat/9708022 (1997);
- [21] T. Sugiyama and N. Nagaosa, Phys. Rev. Lett. **76** (1993) 1980;
- [22] J. Desbois, S. Ouvry and C. Texier, Nucl. Phys B **500**[FS] (1997) 486; Nucl. Phys B **528**[FS] (1998) 727;
- [23] R. Emparan and M.A. Valle Basagoiti, Phys. Rev. B **49** (1994) 14460;
- [24] L. Smrcka and P. Streda, J. Phys. C **10** (1977) 2153;
- [25] Streda, J. Phys. C **15** (1982) L717;
- [26] B.I. Halperin, Phys. Rev. B **25** 2185;
- [27] A.M.M. Pruisken, Nucl. Phys. B **235**[FS11] (1984) 277;
- [28] H. Levine, S.B. Libby and A.M.M. Pruisken, Nucl. Phys. B **240**[FS12] (1984) 30,49,71;
- [29] F. Wegner, Z. Phys. B **35** (1979) 207;
- [30] A.J. McKane and D. Stone, Ann. of Phys. **131** (1981) 36;
- [31] K.B. Efetov, Adv. in Phys. **32** (1983) 53;
- [32] E. Akkermans, A. Auerbach, J.E. Avron and B. Shapiro, Phys. Rev. Lett. **66** (1991) 76;

- [33] J.S. Faulkner, J. Phys. C **10** (1977) 4661;
- [34] A. Moroz, Phys. Rev. A **53** (1996) 669;
- [35] O. Bergman and G. Lozano, Ann. of Phys. **229** (1994) 416;
- [36] S. Ouvry, Phys. Rev. D **50** (1994) 5296;
- [37] R. Jackiw and S.Y. Pi, Phys. Rev. D **42** (1990) 3500;
- [38] A. Comtet, S. Mashkevich and S. Ouvry, Phys. Rev. D **52** (1995) 2594;
- [39] Y. Aharonov and A. Casher, Phys. Rev. A **19** (1979) 2461;
- [40] A.G. Aronov, E. Altshuler, A.D. Mirlin and P. Wölfle, Europhys. Lett. **29** (1995) 239;
- [41] D. Vollhardt and P. Wölfle, Phys. Rev. B **22** (1980) 4666; Phys. Rev. Lett. **48** (1982) 699;
- [42] M. Nielsen and P. Hedegaard, Phys. Rev. B **51** (1995) 7679;
- [43] T. Ando, A.B. Fowler and F. Stern, Rev. Mod. Phys. **54** (1982) 437;